

**NR511 / NR 511 FINAL NEWEST 2024 TEXT BANK
EXAM WITH 1000 VERIFIED ANSWERS GRADED A+
BY EXPERT: DIFFERENTIAL DIAGNOSIS & PRIMARY
CARE PRACTICUM (CHAMBERLAIN)**

Which of the following medications for type 2 diabetes mellitus should not be prescribed during pregnancy?

- Glucotrol

During a DRE on a 75-year-old man, the clinician suspects the patient has prostate cancer. What physical finding should make the clinician suspicious?

- An enlarged rubbery gland

Eddie, age 4, presents to the ED with a live insect trapped in his ear canal causing a lot of distress. What should be your first step?

- Immobilize the insect with 2% lidocaine.

What is the most common cause of microcytic anemia?

- Iron-deficiency anemia

An 82-year-old man is seen in the primary care office with complaints of dribbling urine and difficulty starting his stream. Which of the following should be included in the list of differential diagnoses?

- Benign prostatic hyperplasia (BPH)

Parkinson's disease Prostate cancer

All of the above

Early rheumatoid disease is characterized by:

- Joint swelling and immobility on rising

A 70-year-old female has fallen 2 weeks ago and developed immediate pain in her left wrist. She thought she just bruised it but is worried because it has not improved. She has used Tylenol® and ice at home, and that has helped slightly. You examine her and find she has moderate swelling and ecchymosis but no overtly obvious deformity. Her ROM is uncomfortable and severely diminished due to the pain. No crepitus is heard or felt. Her fingers are warm; her pulse is strong; and capillary refill is less than 2 seconds. What should you do?

- Obtain a wrist x-ray and place her wrist in a splint or prescribe a splint

The criteria for diagnosing generalized anxiety disorder in the American Psychiatric Association's Diagnostic and Statistical Manual of Mental Disorders, 5th edition (text revision) state that excessive worry or apprehension must be present more days than not for at least:

- 6 months

A 60-year-old man presents with an enlarged scrotum. The clinician uses a penlight to transilluminate the scrotum. In a patient with a hydrocele, what would the clinician expect to find? - the scrotum will appear light pink or yellow.

According to Kübler-Ross, the stages of grief occur in which order?

- Denial, anger, bargaining, depression, acceptance

Reuben, age 24, has HIV and just had a routine viral load test done. The results show a falling viral load. What does this indicate?

- A favorable prognostic trend

The exanthem of Lyme disease is - Erythema migrans

You have detected the presence of crepitus on examination of a patient with a musculoskeletal complaint. Additionally, there is limited range of motion (ROM) with both active and passive movement. These findings suggest that the origin of the musculoskeletal complaint is:

- Articular

What is the recommended daily calcium intake for adults over the age of 50 with low bone mass?

- 1,500 mg/day

When assessing a woman for infertility, which of the following tests should be done first?

- Analysis of partner's sperm