

Question: 1 of 180

Time Remaining: 03:59:55
Pause Remaining: 00:05:00

A home health nurse is caring for a child who has Lyme disease. Which of the following is an appropriate action for the nurse to take



- Ensure the state health department has been notified.
- Administer antitoxin.
- Educate the family to avoid sharing personal belongings.
- Assess for skin necrosis.



Question 1 of 180

A home health nurse is caring for a child who has Lyme disease. Which of the following is an appropriate action for the nurse to take?

- Ensure the state health department has been notified.
- Administer antitoxin.
- Educate the family to avoid sharing personal belongings.
- Assess for skin necrosis.

Correct Answer: Ensure the state health department has been notified.

Explanation: Lyme disease is a reportable condition in many states, so the nurse should ensure proper notification of health authorities. Administering antitoxin or assessing for skin necrosis is not applicable, and Lyme disease is not transmitted through shared personal belongings.

Question: 2 of 180

Time Remaining: 03:58:58
Pause Remaining: 00:05:00



A nurse is caring for a client who has been admitted to the hospital.

Exhibit 1 Exhibit 2 Exhibit 3

Laboratory Results

1200:

Hgb 9.5 g/dL (14 to 18 g/dL)

Hct 38% (42% to 52%)

Bilirubin 5.3 mg/dL (0.3 to 1.0 mg/dL)

Creatinine 1.8 mg/dL (0.6 to 1.3 mg/dL)

Platelet count 100,000/mm³ (150,000 to 400,000/mm³)

1800:

Alanine aminotransferase ALT 51 units/L (4 to 36 units/L)

Aspartate aminotransferase AST 48 units/L (0 to 35 units/L)

Alkaline phosphate ALP 151 units/L (30 to 120 units/L)

Blood total protein 15 g/dL (6.4 to 8.3 g/dL)

Select the 5 actions the nurse should take.



- Provide frequent rest periods for the client.
- Restrict the client's sodium intake.
- Advise the client to avoid the use of soap and alcohol-based lotions.
- Place the client on a low-carbohydrate diet.
- Place the client under contact isolation.
- Instruct the client to avoid blowing their nose forcefully.
- Assess the client's level of orientation.

Question: 3 of 180

Time Remaining: 03:56:10
Pause Remaining: 00:05:00

PAUSE



A nurse is caring for a client who has a vented NG tube set to low intermittent suction and has vomited. Which of the following actions should the nurse perform first?



- Administer an antiemetic medication.
- Evaluate functioning of the suction device.
- Provide oral hygiene care.
- Replace the NG tube.

Question: 4 of 180

Time Remaining: 03:55:49
Pause Remaining: 00:05:00

PAUSE



While performing a routine assessment, a nurse notices fraying on the electrical cord of a client's continuous passive motion (CPM) device. Which of the following actions should the nurse take first?

- Initiate a requisition for a replacement CPM device.
- Report the defect to the equipment maintenance staff.
- Remove the device from the room.
- Ensure the device inspection sticker is current.

Question: 5 of 180

Time Remaining: 03:53:46
Pause Remaining: 00:05:00

PAUSE



A nurse is setting up a sterile field to perform wound irrigation for a client. Which of the following actions should the nurse take when pouring the sterile solution?

- Remove the cap and place it sterile-side up on a clean surface.
- Place sterile gauze over areas of spilled solution within the sterile field.
- Hold the bottle in the center of the sterile field when pouring the solution.
- Hold the irrigation solution bottle with the label facing away from the palm of the hand.