

# ATI COMPREHENSIVE EXIT EXAM 2023 WITH NGN

Question: 1 of 180

Time Remaining: 03:59:55  
Pause Remaining: 00:05:00

A home health nurse is caring for a child who has Lyme disease. Which of the following is an appropriate action for the nurse to take?

- Ensure the state health department has been notified.
- Administer antitoxin.
- Educate the family to avoid sharing personal belongings.
- Assess for skin necrosis.

Question: 2 of 180

Time Remaining: 03:58:58  
Pause Remaining: 00:05:00

PAUSE



A nurse is caring for a client who has been admitted to the hospital.

Exhibit 1

Exhibit 2

Exhibit 3



### Laboratory Results

1200:

Hgb 9.5 g/dL (14 to 18 g/dL)

Hct 38% (42% to 52%)

Bilirubin 5.3 mg/dL (0.3 to 1.0 mg/dL)

Creatinine 1.8 mg/dL (0.6 to 1.3 mg/dL)

Platelet count 100,000/mm<sup>3</sup> (150,000 to 400,000/mm<sup>3</sup>)

1800:

Alanine aminotransferase ALT 51 units/L (4 to 36 units/L)

Aspartate aminotransferase AST 48 units/L (0 to 35 units/L)

Alkaline phosphate ALP 151 units/L (30 to 120 units/L)

Blood total protein 15 g/dL (6.4 to 8.3 g/dL)

Select the 5 actions the nurse should take.

- Provide frequent rest periods for the client.
- Restrict the client's sodium intake.
- Advise the client to avoid the use of soap and alcohol-based lotions.
- Place the client on a low-carbohydrate diet.
- Place the client under contact isolation.
- Instruct the client to avoid blowing their nose forcefully.
- Assess the client's level of orientation.

Question: 3 of 180

Time Remaining: 03:56:10  
Pause Remaining: 00:05:00

PAUSE



A nurse is caring for a client who has a vented NG tube set to low intermittent suction and has vomited. Which of the following actions should the nurse perform first?



- Administer an antiemetic medication.
- Evaluate functioning of the suction device.
- Provide oral hygiene care.
- Replace the NG tube.

Question: 4 of 180

Time Remaining: 03:55:49  
Pause Remaining: 00:05:00

PAUSE



While performing a routine assessment, a nurse notices fraying on the electrical cord of a client's continuous passive motion (CPM) device. Which of the following actions should the nurse take first?

- Initiate a requisition for a replacement CPM device.
- Report the defect to the equipment maintenance staff.
- Remove the device from the room.
- Ensure the device inspection sticker is current.

Question: 5 of 180

Time Remaining: 03:53:46  
Pause Remaining: 00:05:00

PAUSE



A nurse is setting up a sterile field to perform wound irrigation for a client. Which of the following actions should the nurse take when pouring the sterile solution?

- Remove the cap and place it sterile-side up on a clean surface.
- Place sterile gauze over areas of spilled solution within the sterile field.
- Hold the bottle in the center of the sterile field when pouring the solution.
- Hold the irrigation solution bottle with the label facing away from the palm of the hand.

Question: 6 of 180

Time Remaining: 03:53:03  
Pause Remaining: 00:05:00

PAUSE



A nurse is creating a plan of care for a female client who has recurrent urinary tract infections. Which of the following interventions should the nurse include in the plan?



- Wear loose-fitting underwear.
- Take a bubble bath after intercourse.
- Drink four 240 mL (8 oz) glasses of water each day.
- Void every 5 to 6 hr during the day.

Question: 7 of 180

Time Remaining: 03:52:35  
Pause Remaining: 00:05:00

PAUSE



A nurse is caring for a newborn.

Exhibit 1   Exhibit 2   Exhibit 3

Vital Signs

0640:

Temperature 36.7° C (98.1° F) axillary  
Heart rate 154/min  
Respiratory rate 68/min  
BP 72/48 mm Hg


0650:

Heart rate 156/min  
Respiratory rate 72/min

0700:

Temperature 37° C (98.6° F) axillary  
Heart rate 156/min  
Respiratory rate 76/min

Drag words from the choices below to fill in each blank in the following sentence.



The client is at risk for developing Target 1 and

Target 2

**Findings**

- hypoglycemia
- bronchopulmonary dysplasia
- transient tachypnea of the newborn
- tachycardia