

# Question: 1 of 70

Time Remaining: 01:28:13  
Pause Remaining: 00:05:00

PAUSE



A nurse is caring for a client who is experiencing a postpartum hemorrhage and has a new prescription for misoprostol.

Exhibit 1

Exhibit 2

Exhibit 3

Exhibit 4

## Medical History

Preeclampsia

Cesarean birth of viable twin male newborns

The nurse is assessing the client 30 min later. How should the nurse interpret the findings?

For each finding, click to specify whether the finding is unrelated to the diagnosis, an indication of potential improvement, or an indication of potential worsening condition.

Findings 30 min later	Unrelated to diagnosis	Indication of potential improvement	Indication of potential worsening condition
Fundus at level of umbilicus	<input type="radio"/>	<input checked="" type="radio"/>	<input type="radio"/>
Cloudy urine	<input type="radio"/>	<input type="radio"/>	<input checked="" type="radio"/>
Blood pressure 80/50 mm Hg	<input type="radio"/>	<input type="radio"/>	<input checked="" type="radio"/>
Moderate lochia rubra	<input type="radio"/>	<input checked="" type="radio"/>	<input type="radio"/>
Thready pulse	<input type="radio"/>	<input type="radio"/>	<input checked="" type="radio"/>
Fundus firm to palpation	<input type="radio"/>	<input checked="" type="radio"/>	<input type="radio"/>

CONTINUE

A nurse is assessing a postpartum client during a follow-up visit.

- Exhibit 1
- Exhibit 2
- Exhibit 3**

**Vital Signs**

0930:  
Temperature 37° C (98.6° F)  
Pulse rate 78/min  
Respiratory rate 12/min  
Blood pressure 124/80 mm Hg  
Pulse oximetry 100%

Complete the following sentence by using the list of options.



The nurse is teaching the client about postpartum depression. The nurse should encourage the client to maintain a set schedule and exercise 30 min per day to help prevent postpartum depression.



CONTINUE



Full screen mode is in effect during your proctored testing.  
Please Note: Attempting to exit out of full screen mode or pressing the escape button will result in an alert message sent to the proctor and may result in the termination of the assessment.

## Question: 3 of 70

Time Remaining  
Pause Remaining

A nurse is assessing a postpartum client who delivered vaginally 8 hr ago.

Exhibit 1

Exhibit 2

Nurses' Notes

0700:

Breasts soft, nipples intact. Uterus palpated firm, midline, and at level of umbilicus. Moderate amount of lochia rubra. Episiotomy site well approximated with mild edema and ecchymosis. Client reports pain as 2 on a scale of 0 to 10. Able to void spontaneously; no bladder distention. Deep tendon reflexes 1+. Peripheral edema 2+ in bilateral lower extremities.

1100:

Breasts soft, nipples intact. Uterus palpated soft with lateral deviation and 1 cm above the umbilicus. Large amount of lochia rubra. Episiotomy site well approximated with mild edema and ecchymosis. Client reports pain as 3 on a scale of 0 to 10. Deep tendon reflexes 1+. Peripheral edema 2+ in bilateral lower extremities.

Select the 3 findings that require immediate follow-



- Lateral deviation of the uterus
- Deep tendon reflexes 1+
- Peripheral edema 2+ bilateral lower extremities
- Pain rating of 3 on a scale of 0 to 10 (increased)
- Uterine tone soft
- Large amount of lochia rubra
- Blood pressure 136/86 mm Hg
- Breasts soft



## Question: 5 of 70

Time Remaining: 01:25:01

Pause Remaining: 00:05:00

PAUSE



A nurse is caring for a client who is receiving an epidural block with an opioid analgesic. The nurse should monitor for which of the following findings as an adverse effect of the medication?



Hyperglycemia

Bilateral crackles

P 84

Hypotension

Polyuria