

Question 3

Time remaining: 00:00:00

While performing a routine assessment, a nurse notices fraying on the electrical cord of a client's continuous passive motion (CPM) device. Which of the following actions should the nurse take first?



- Initiate a requisition for a replacement CPM device.
- Report the defect to the equipment maintenance staff.
- Remove the device from the room.
- Ensure the device inspection sticker is current.

Question 4

Time remaining: 00:00:00

A nurse is setting up a sterile field to perform wound irrigation for a client. Which of the following actions should the nurse take when pouring the sterile solution?



- Remove the cap and place it sterile-side up on a clean surface.
- Place sterile gauze over areas of spilled solution within the sterile field.
- Hold the bottle in the center of the sterile field when pouring the solution.
- Hold the irrigation solution bottle with the label facing away from the palm of the hand.

Question 5

Time remaining: 00:00:00

A nurse is creating a plan of care for a female client who has recurrent urinary tract infections. Which of the following interventions should the nurse include in the plan?



- Wear loose-fitting underwear.
- Take a bubble bath after intercourse.
- Drink four 240 mL (8 oz) glasses of water each day.
- Void every 5 to 6 hr during the day.

Question 6

Time remaining: 00:05:00

A nurse is conducting health promotion education regarding contraindications to combination oral contraceptive use to a group of women. Which of the following conditions should the nurse include in the teaching?



- Hypertension
- Fibromyalgia
- Renal calculi
- Fibrocystic breast disease

Question 7

Time remaining: 00:05:00

A nurse is caring for a client who is immobile. Which of the following interventions is appropriate to prevent contracture?



- Position a pillow under the client's knees.
- Place a towel roll under the client's neck.
- Align a trochanter wedge between the client's legs.
- Apply an orthotic to the client's foot.

Question 8

Time remaining: 00:05:00

A nurse is caring for a client who is experiencing expressive aphasia and right hemiparesis following a stroke. Which of the following actions by the nurse best promotes communication among staff caring for the client?



- Posting swallowing precautions at the head of the client's bed
- Noting changes in the treatment plan in the client's medical record
- Recording the client's progress in the nurses' notes
- Having interdisciplinary team meetings for the client on a regular basis

Question 9

Pause Remaining: 00:05:00



A nurse has just received change-of-shift report for four clients. Which of the following clients should the nurse assess first?



- A client who is scheduled for a procedure in 1 hr
- A client who received a pain medication 30 min ago for postoperative pain
- A client who was just given a glass of orange juice for a low blood glucose level
- A client who has 100 mL of fluid remaining in his IV bag

CONTINUE

Question 10

A nurse is performing postmortem care for a recently deceased client prior to the client's family visit. Which of the following actions should the nurse plan to take?



- Cross the client's arms across their chest.
- Hold the client's eyes shut for a few seconds.
- Place the client in a high-Fowler's position.
- Remove the client's dentures from their mouth.

Question 11

A nurse is caring for a client who has been admitted to the hospital.

Exhibit 1

Exhibit 2

Exhibit 3

Laboratory Results

1200:

Hgb 9.5 g/dL (14 to 18 g/dL)
Hct 38% (42% to 52%)
Bilirubin 5.3 mg/dL (0.3 to 1.0 mg/dL)
Creatinine 1.8 mg/dL (0.6 to 1.3 mg/dL)
Platelet count 100,000/mm³ (150,000 to 400,000/mm³)

1800:

Alanine aminotransferase ALT 51 units/L (4 to 36 units/L)
Aspartate aminotransferase AST 48 units/L (0 to 35 units/L)
Alkaline phosphate ALP 151 units/L (30 to 120 units/L)
Blood total protein 15 g/dL (6.4 to 8.3 g/dL)

Select the 5 actions the nurse should take.



- Provide frequent rest periods for the client.
- Restrict the client's sodium intake.
- Advise the client to avoid the use of soap and alcohol-based lotions.
- Place the client on a low-carbohydrate diet.
- Place the client under contact isolation.
- Instruct the client to avoid blowing their nose forcefully.
- Assess the client's level of orientation.