ATI COMPREHENSIVE EXITV2 FINAL2024Targeted Exam EXAM LATEST WITH ACTUAL QUESTIONS AND CORRECT VERIFIED ANSWERS ALREADY GRADED A+ 100% GUARANTEED PASS!

Amitriptyline adverse reaction to monitor for -

√√ANSWER√√>>urinary retention

Group therapy for families adapting to loss of family members is -

√√ANSWER√√>>determine roles

First step to perform when a patient says they want to have a family

home visit - $\sqrt{ANSWER}\sqrt{>>}$ clarify the source

Referral that should be made for a patient who has a non-healing

ulcer - √√ANSWER√√>>dietitian referral

Home safety for an older adult - \sqrt{ANSWER} , $\sqrt{>}$ > install grab bars in

the tub

Patient has a potassium level of 5.2, what med should be held? -

√√ANSWER√√>>spironolactone (because its potassium-sparing)

Patient states they wish they were dead, nurse should respond by

saying - \sqrt{ANSWER} , $\sqrt{>}$ "you seem to be feeling hopeless"

Acute MI with alteplase contraindication -

cardiac failure 6 months ago

To promote sleep - √√ANSWER√√>>eat a small bedtime snack

Home tracheostomy care, correct statement by family is -

 \checkmark ANSWER \checkmark >>have suction set to 60 mm Hg

Complication of immobility - $\sqrt{ANSWER}\sqrt{>}$ swollen area in calf

Phenelzine - √√ANSWER√√>>smoked fish

Interventions to decrease cardiac demands -

√√ANSWER√√>>Maintain bed rest

Position at 45 degree angle- keep safety restraints low & loose on

abdomen

Allow child to sleep with several pillows & encourage semi fowlers or

fowlers position while awake

Feeding techniques for cleft palate - √√ANSWER√√>>Position

infant upright while cradling the head

Use specialized bottle with one way valve & special cut nipple

Burp frequently

Neuroleptic malignant syndrome - \sqrt{ANSWER} ,

increased BP, muscle rigidity

Correct statement for patient with stomatitis -\sqrt{ANSWER}

a soft bristle toothbrush

Decrease cardiac demands in an infant you should -

√√ANSWER√√>>Feed the infant when awake & crying

Cocaine patient s/s - √√ANSWER√√>>Increased temperature

Instruction for the removal of an NG tube - \sqrt{ANSWER}

deep breath & hold it

With a continuous NG feeding, you should -

 \sqrt{ANSWER} , $\sqrt{>}$ >Suction gas residuals every 4 hours

Patient who is most at risk for aspirating - \sqrt{ANSWER}

a history of GERD

Expected finding for retinoblastoma - \sqrt{ANSWER}

reflex

Teaching for the parents of a child who is taking Digoxin -

 \sqrt{ANSWER}

Patient who is at risk for developing streptococcus -

 \sqrt{ANSWER} , $\sqrt{>>7}$ year old recovering from impetigo

Procedure that requires sterile gloves -

√√**ANSWER**√**√**>>Tracheostomy care