

ATI PN COMPREHENSIVE EXIT EXAM 2024/2025 WITH (ACTUAL EXAM)QUESTIONS AND 100% VERIFIED ANSWERS | GRADED A+

BUN levels - ANSWERS-10-20

WBC levels - ANSWERS-5,000-10,000

Serum creatinine levels - ANSWERS-males 0.6-1.2; females 0.5-1.1

Therapeutic digoxin levels - ANSWERS-0.8-2.0

HbA1c levels - ANSWERS-4%-6%

Fasting glucose levels - ANSWERS-70-105

HDL levels - ANSWERS-males 35-65; females 35-85

Total serum cholesterol levels - ANSWERS-less than 200

Tb

Family precautions

How to get cleared - ANSWERS-Family already exposed

3 negative cultures required.

comminuted fracture - ANSWERS-fracture in which the bone is splintered or crushed

Most problems of alkalosis are related to increased stimulation of which three systems - ANSWERS-cardiac,

nervous

neuromuscular systems

The second stage of labor begins when what two things happen - [ANSWERS](#)-cervix is completely dilated and there is a strong desire to push

Fetus HR when compressed by birth canal - [ANSWERS](#)-HR slows

aphasic - [ANSWERS](#)-without speech

When and how should a Breast exam be done - [ANSWERS](#)-7 days after menstrual cycle with pads after

The normal clotting time is - [ANSWERS](#)-8 to 15 minutes.

The normal ammonia value - [ANSWERS](#)-10 to 80 mcg/dL

therapeutic range for a serum phenytoin level is - [ANSWERS](#)-10 to 20 mcg/mL

Guillain-Barré common systems affected - [ANSWERS](#)-resp and issues

Eye drop and ointment sequence when prescribed together - [ANSWERS](#)-Eye drop should be followed by ointment separated by 3-5 min

High temp after delivery means - [ANSWERS](#)-dehydration

Most common feeling at the end of life - [ANSWERS](#)-Anxiety most common at end of life

Calcium carbonate is best administered - [ANSWERS](#)-1 to 1½ hours after meals with full glass of water

Myringotomy - ANSWERS-incision into the tympanic membrane

Cystitis - ANSWERS-inflammation of the bladder

Regression of behaviors in a toddler when sick - ANSWERS-is normal in toddlers when stressed. To be ignored

What time of day is IOP higher - ANSWERS-IOP is higher in the mornings

frequency to check IV - ANSWERS- IV's should be checked every hour

When can a hep V patient bottle feed - ANSWERS-Hep B may bottle feed after vaccination

Nitro

How long is it good after opening

Indications - ANSWERS-Nitro is good for 6 months

may be used protractedly or for the onset of angina

Salt values

Less adrenals

Heart failure

Cushions - ANSWERS-Less adrenals less salt.

Heart failure and Cushing more salt.

Prolapsed Cord: Care - ANSWERS-Care Includes: Knee to Chest position OR Trendelenburg

Keep it from drying and shrinking by maintaining moisture

Dietary Recommendations for nausea - ANSWERS-Cool clear liquids help nausea

platypelloid pelvis - ANSWERS-pelvis that is flat in its dimensions with a very narrow anterior-posterior diameter and a wide transverse diameter; this shape makes it extremely difficult for the fetus to pass through the bony pelvis

dystocia - ANSWERS-difficult labor

Sodium equilibrium with potassium - ANSWERS-Sodium exchanges with potassium

Pregnancy prediction - ANSWERS-Subtract 3 months on the same day,
add 7 days
then add a FULL year.

Alendronate sodium (Fosamax) is for Osteoporosis. How should it be taken - ANSWERS-taken on empty stomach with water only or useless.

Vascular headaches treated and side effects - ANSWERS-with meds that cause vasoconstriction.

Cool numb extremities and hypertension as result

Quickening or fetal movement is felt - ANSWERS-at 16-20 wks.

Stockings application timing - ANSWERS-elevate legs 15 to 30 min prior to applying the elastic stockings to decrease swelling and improve circulation to his legs

How to determine the client's actual urinary output during irrigation - ANSWERS-record the amount of fluid used for irrigation and the amount returned.

Post TURP flow rate - ANSWERS-maintain the flow rate of the bladder irrigation to keep the urine diluted to a reddish-pink color and the tubing free of clots and bleeding

If tube becomes clogged - ANSWERS-flush the feeding tube with 30 to 60 mL of warm water