ATI RN MATERNAL NEWBORN 2024 PROCTOR EXAM WITH NGN LATEST VERSION WITH 70 QUESTIONS.

NURSINGONLINEEXAMS@GMAIL.COM

QUESTION 1

A nurse is caring for a client who has been admitted to the antepartum Click to highlight the findings that require follow-up. To deselect a finding, click on the finding again. unit. Exhibit 1 Exhibit 2 Exhibit 3 **History and Physical Vital Signs** Day 1, 0900: Day 1, 0900: 30-year-old client at 33 weeks gestation, Gravida 4 Para 3 Admission: Maternal blood type: Rh+ Temperature 38.4° C (101.1° F) Last pregnancy resulted in a preterm spontaneous vaginal birth at 30 weeks Heart rate 92/min gestation. Respiratory rate 18/min Blood pressure 130/78 mm Hg Nurses' Notes Pre-pregnancy BMI 27.6 Day 1, 0900: Current BMI 29.9 Client reports lower back pain and pinkish vaginal discharge. Uterine contractions every 8 minutes, palpate strong, duration 30 seconds. FHR baseline 145, minimal variability. Cervical exam indicates 2 cm, 50% effaced, 0 station. CBC and urinalysis collected and sent to lab.

QUESTION 2

A nurse is caring for a client who has been admitted to the antepartum

Exhibit 1 Exhibit 3

History and Physical

Day 1, 0900:

30-year-old client at 33 weeks gestation, Gravida 4 Para 3 Maternal blood type: Rh+

Last pregnancy resulted in a preterm spontaneous vaginal birth at 30 weeks gestation.

NKA

Click to highlight the findings that require follow-up. To deselect a finding, click on the finding again.

History and Physical

Day 1, 0900:

30-year-old client at 33 weeks gestation, Gravida 4 Para 3

Maternal blood type: Rh+

Last pregnancy resulted in a preterm spontaneous vaginal birth at 30 weeks gestation.

Nurses' Notes

Day 1, 0900:

Client reports lower back pain and pinkish vaginal discharge.

Uterine contractions every 8 minutes, palpate strong, duration 30 seconds.


FHR baseline 145, minimal variability.

Cervical exam indicates 2 cm, 50% effaced, 0 station.

Membranes intact.

CBC and urinalysis collected and sent to lab.

QUESTION 3



QUESTION 4

A nui					_			7			abo	out	br	ea:	tfe	eed	ling	g h	er	2-v	vee	ek-	old	in	fan	t. V	Vhi	ch	of	the	fo	llo	win	g s	tat	em	en	ts	by	the	e pa	rei	nt	ind	lica	ites	an			
		0 4	•						0				•							•	0				•	0		•			0		•		٠	•		•		•	•	0	•		•	0		•	•	٠
0)	"Afi	ter	5 to	10) mi	nut	es	whe	en	the	br	ea	st is	er	np	tiec	i, n	ny l	oat	oy s	sho	uld	i be	e re	mo	ve	d fr	om	th	e b	rea	st."																	
0)	"Ma	anı	ually	/ ex	pre	ssir	ng r	ny i	mil	k v	vill	de	crea	se	m	y m	ilk	su	opl	y."																													
)	"My	y b	aby	she	oulc	l alv	way	s si	tari	to	n th	ne s	san	ne b	ore	ast	wł	nen	fe	edi	ing	."																											
0)	"Th	ne n	nor	e m	y b	aby	is a	at tl	ne	bre	eas	t sı	uck	ng	, th	e n	nor	e n	nill	clv	vill	pro	odu	ice.	**																								

Q	UESTION 5
	A nurse is providing teaching about immunizations to a client who is pregnant. Which of the following statements should the nurse include teaching?
	"The immunization for varicella should be given at least 1 month prior to delivery."
	"You can receive the rubella immunization during the third trimester of pregnancy."
	"The hepatitis B immunization should not be obtained until after you finish breastfeeding."
ι	"You can receive the immunization for influenza at any time during your pregnancy." JESTION 6
	WESTION 6 A nurse is caring for a client who is 12 hr postpartum and has a third-degree perineal laceration. The client reports not having a bowel move for 4 days. Which of the following medications should the nurse administer?
	WESTION 6 A nurse is caring for a client who is 12 hr postpartum and has a third-degree perineal laceration. The client reports not having a bowel move for 4 days. Which of the following medications should the nurse administer?
	WESTION 6 A nurse is caring for a client who is 12 hr postpartum and has a third-degree perineal laceration. The client reports not having a bowel move for 4 days. Which of the following medications should the nurse administer?
	WESTION 6 A nurse is caring for a client who is 12 hr postpartum and has a third-degree perineal laceration. The client reports not having a bowel move for 4 days. Which of the following medications should the nurse administer? Bisacodyl 10 mg rectal suppository Magnesium hydroxide 30 mL PO
	WESTION 6 A nurse is caring for a client who is 12 hr postpartum and has a third-degree perineal laceration. The client reports not having a bowel move for 4 days. Which of the following medications should the nurse administer?

A nurse is developing a plan of care for a newborn who is to undergo phototherapy for hyperbillirubinemia. Which of the following actions should the nurse include in the plan?
- ✓✓ANSWER✓✓>>Remove all clothing form the newborn except the diaper

A nurse is caring for a client who is at 15 weeks gestation, is Rh-negative, and has just had an amniocentesis. Which of the following interventions is the nurse's priority following the procedure? - \(\sqrt{ANSWER} \sqrt{V} >> \) Monitor the FHR

A nurse in an antepartum clinic is assessing a client who is at 32 weeks of gestation. Which of the following findings should the nurse report to the provider? ✓✓ANSWER✓✓>>Reports of decreased fetal movement

A nurse is caring for a client who is at 36 weeks of gestation and has a positive contraction stress test. The nurse should plan to prepare the client for which of the following diagnostic test? - \sqrt{A} ANSWER \sqrt{A} >>Biophysicial profile

A nurse is preparing to administer magnesium sulfate 2g/hr IV to a client who is in preterm labor. Available is 20 g magnesium sulfate in 500 mL of dextrose 5% in water (D5W). The nurse should set the IV infusion pump to administer how many mL/hr? - $\sqrt{ANSWER}\sqrt{J}>>50$ ml/hr

A nurse is caring for a client who has hyperemesis gravidarum and is receiving IV fluid replacement. Which of the following findings should the nurse report to the provider? - VANSWERV>>BUN 25 mg/dL

A nurse is assessing a late preterm newborn. Which of the following manifestations is an indication of hypoglycemia? - $\sqrt{ANSWER}\sqrt{>}>Respiratory$ distress

A nurse in an antepartum clinic is providing care for a client who is at 26 weeks of gestation. Upon reviewing the client's medical record, which of the following findings should the nurse report to the provider? - \(\sqrt{ANSWER} \sqrt{I} >> \) Fundal Height Measurement